

2016-10-07 Exam in

Turbulence modeling, MTF270

- **Time:** 14.00–18.00 **Location:** V
 - **Teacher:** Lars Davidson, phone 772 1404, 0730-791 161
 - **Aids**
 - Formula sheet, 2 pages, appended to the exam
 - **Checking the evaluation and results of your written exam:** At 12-13: Oct 31 and Nov 1 in Room Euler
 - **Grading:** 20-29p: 3, 30-39: 4, 40-50: 5.
-

T1. a) How is the buoyancy term, ρg_i , re-written in incompressible flow? (5p)

b) How is the production term modeled in the $k - \varepsilon$ model? Show how it can be expressed in \bar{s}_{ij} (5p)

T2. a) The exact Poisson equation for the pressure fluctuation reads (5p)

$$\frac{1}{\rho} \frac{\partial^2 p'}{\partial x_j \partial x_j} = -2 \underbrace{\frac{\partial \bar{v}_i}{\partial x_j} \frac{\partial v'_j}{\partial x_i}}_{\text{rapid term}} - \underbrace{\frac{\partial^2}{\partial x_i \partial x_j} (v'_i v'_j - \overline{v'_i v'_j})}_{\text{slow term}} \quad (1)$$

Derive this equation.

b) Consider streamline curvature for a streamline formed as a circular arc (convex curvature). Show that the turbulence is dampened if $\partial v_\theta / \partial r > 0$ and that it is enhanced if the sign of $\partial v_\theta / \partial r$ is negative. (5p)

T3. a) What is a realizability constraint? Give the main realizability constraint for the normal stress and the shear stress. The Boussinesq assumption may violate one of those two constraints: show which one. (5p)

b) Show how a sinus wave $\sin(\kappa_c x)$ corresponding to cut-off is represented on a grid with two and four nodes, respectively. How is κ_c related to the grid size Δx for these cases? (5p)

- T4. a) Consider the energy spectrum. Show the three different regions (the large energy-containing scales, the $-5/3$ range and the dissipating scales). In which region should the cut-off be located? How are k , k_{res} and k_{sgs} computed from the energy spectrum? (5p)
- b) Consider a 1D finite volume grid. Carry out a second filtering of \bar{v} at node I and show that $\bar{\bar{v}}_I \neq \bar{v}_I$. (5p)
- T5. a) Explain what is DES. The length scale in the RANS S-A model reads $\left(\frac{\tilde{\nu}_t}{d}\right)^2$; how is it computed in the corresponding DES model? (5p)
- b) Consider the SAS model. How is the von Kármán length scale defined? An additional source term is introduced in the ω equation: what is the form of this term? What is the object of this term? When is it large and small, respectively? (5p)

MTF270 Turbulence modeling: Formula sheet

October 5, 2016

The continuity, Navier-Stokes and temperature equations for incompressible flow with constant viscosity read (*conservative form*)

$$\begin{aligned}\frac{\partial v_i}{\partial x_i} &= 0 \\ \rho_0 \frac{\partial v_i}{\partial t} + \rho_0 \frac{\partial v_i v_j}{\partial x_j} &= -\frac{\partial p}{\partial x_i} + \mu \frac{\partial^2 v_i}{\partial x_j \partial x_j} - \rho_0 \beta (\theta - \theta_0) g_i \\ \frac{\partial \theta}{\partial t} + \frac{\partial v_i \theta}{\partial x_i} &= \alpha \frac{\partial^2 \theta}{\partial x_i \partial x_i}\end{aligned}$$

► The Navier-Stokes equation for incompressible flow with constant viscosity read (*non-conservative form*, p denotes the hydrostatic pressure, i.e. $p = 0$ if $v_i = 0$)

$$\rho_0 \frac{\partial v_i}{\partial t} + \rho_0 v_j \frac{\partial v_i}{\partial x_j} = -\frac{\partial p}{\partial x_i} + \mu \frac{\partial^2 v_i}{\partial x_j \partial x_j}$$

The *time averaged* continuity equation, Navier-Stokes equation temperature equations read

$$\begin{aligned}\frac{\partial \bar{v}_i}{\partial x_i} &= 0 \\ \rho_0 \frac{\partial \bar{v}_i \bar{v}_j}{\partial x_j} &= -\frac{\partial \bar{p}}{\partial x_i} + \frac{\partial}{\partial x_j} \left(\mu \frac{\partial \bar{v}_i}{\partial x_j} - \rho_0 \overline{v'_i v'_j} \right) - \rho_0 \beta (\bar{\theta} - \theta_0) g_i \\ \frac{\partial \bar{v}_i \bar{\theta}}{\partial x_i} &= \alpha \frac{\partial^2 \bar{\theta}}{\partial x_i \partial x_i} - \frac{\partial \overline{v'_i \theta'}}{\partial x_i}\end{aligned}$$

The Boussinesq assumption reads

$$\overline{v'_i v'_j} = -\nu_t \left(\frac{\partial \bar{v}_i}{\partial x_j} + \frac{\partial \bar{v}_j}{\partial x_i} \right) + \frac{2}{3} \delta_{ij} k = -2\nu_t \bar{s}_{ij} + \frac{2}{3} \delta_{ij} k$$

The modeled $\overline{v'_i v'_j}$ equation with IP model reads

$$\begin{aligned}\bar{v}_k \frac{\partial \overline{v'_i v'_j}}{\partial x_k} &= \text{(convection)} \\ -\overline{v'_i v'_k} \frac{\partial \bar{v}_j}{\partial x_k} - \overline{v'_j v'_k} \frac{\partial \bar{v}_i}{\partial x_k} &= \text{(production)} \\ -c_1 \frac{\varepsilon}{k} \left(\overline{v'_i v'_j} - \frac{2}{3} \delta_{ij} k \right) &= \text{(slow part)} \\ -c_2 \left(P_{ij} - \frac{2}{3} \delta_{ij} P^k \right) &= \text{(rapid part)} \\ +c_{1w} \rho_0 \frac{\varepsilon}{k} \left[\overline{v'_k v'_m} n_k n_m \delta_{ij} - \frac{3}{2} \overline{v'_i v'_k} n_k n_j \right. \\ &\quad \left. - \frac{3}{2} \overline{v'_j v'_k} n_k n_i \right] f \left[\frac{\ell_t}{x_n} \right] &= \text{(wall, slow part)} \\ +c_{2w} \left[\Phi_{km,2} n_k n_m \delta_{ij} - \frac{3}{2} \Phi_{ik,2} n_k n_j \right. \\ &\quad \left. - \frac{3}{2} \Phi_{jk,2} n_k n_i \right] f \left[\frac{\ell_t}{x_n} \right] &= \text{(wall, rapid part)} \\ +\nu \frac{\partial^2 \overline{v'_i v'_j}}{\partial x_k \partial x_k} &= \text{(viscous diffusion)} \\ +\frac{\partial}{\partial x_k} \left[c_k \overline{v'_k v'_m} \frac{k}{\varepsilon} \frac{\partial \overline{v'_i v'_j}}{\partial x_m} \right] &= \text{(turbulent diffusion)} \\ -g_i \beta \overline{v'_j \theta'} - g_j \beta \overline{v'_i \theta'} &= \text{(buoyancy production)} \\ -\frac{2}{3} \varepsilon \delta_{ij} &= \text{(dissipation)}\end{aligned}$$

► The exact transport equation for turbulent heat flux vector $\overline{v'_i \theta'}$ reads

$$\begin{aligned} \frac{\partial \overline{v'_i \theta'}}{\partial t} + \frac{\partial}{\partial x_k} \overline{v_k v'_i \theta'} &= \underbrace{-\overline{v'_i v'_k} \frac{\partial \bar{\theta}}{\partial x_k}}_{P_{i\theta}} - \underbrace{\overline{v'_k \theta'} \frac{\partial \bar{v}_i}{\partial x_k}}_{\Pi_{i\theta}} - \underbrace{\frac{\bar{\theta}'}{\rho} \frac{\partial p'}{\partial x_i}}_{\Pi_{i\theta}} - \underbrace{\frac{\partial}{\partial x_k} \overline{v'_k v'_i \theta'}}_{D_{i\theta,t}} \\ &+ \underbrace{\alpha \frac{\partial}{\partial x_k} \left(\overline{v'_i \frac{\partial \theta'}{\partial x_k}} \right)}_{D_{i\theta,\nu}} + \underbrace{\nu \frac{\partial}{\partial x_k} \left(\overline{\theta' \frac{\partial v'_i}{\partial x_k}} \right)}_{\varepsilon_{i\theta}} - (\nu + \alpha) \underbrace{\frac{\partial v'_i}{\partial x_k} \frac{\partial \theta'}{\partial x_k}}_{\varepsilon_{i\theta}} - \underbrace{g_i \beta \overline{\theta'^2}}_{G_{i\theta}} \end{aligned}$$

► The exact k equation reads

$$\frac{\partial k}{\partial t} + \frac{\partial \bar{v}_j k}{\partial x_j} = -\overline{v'_j v'_j} \frac{\partial \bar{v}_i}{\partial x_j} - \frac{\partial}{\partial x_j} \left[\frac{1}{\rho} \overline{v'_j p'} + \frac{1}{2} \overline{v'_j v'_i v'_i} - \nu \frac{\partial k}{\partial x_j} \right] - \nu \frac{\partial v'_i}{\partial x_j} \frac{\partial v'_i}{\partial x_j}$$

► The exact $\overline{v'_i v'_j}$ equation reads

$$\begin{aligned} \frac{\partial \overline{v'_i v'_j}}{\partial t} + \frac{\partial}{\partial x_k} (\overline{v_k v'_i v'_j}) &= -\overline{v'_j v'_k} \frac{\partial \bar{v}_i}{\partial x_k} - \overline{v'_i v'_k} \frac{\partial \bar{v}_j}{\partial x_k} \\ &- \frac{\partial}{\partial x_k} \left(\overline{v'_i v'_j v'_k} + \frac{1}{\rho} \delta_{jk} \overline{v'_i p'} + \frac{1}{\rho} \delta_{ik} \overline{v'_j p'} - \nu \frac{\partial v'_i v'_j}{\partial x_k} \right) \\ &+ \frac{1}{\rho} p' \left(\frac{\partial v'_i}{\partial x_j} + \frac{\partial v'_j}{\partial x_i} \right) - g_i \beta \overline{v'_j \theta'} - g_j \beta \overline{v'_i \theta'} - 2\nu \frac{\partial v'_i}{\partial x_k} \frac{\partial v'_j}{\partial x_k} \end{aligned}$$

► The modeled k and ε equations

$$\begin{aligned} \frac{\partial k}{\partial t} + \bar{v}_j \frac{\partial k}{\partial x_j} &= \nu_t \left(\frac{\partial \bar{v}_i}{\partial x_j} + \frac{\partial \bar{v}_j}{\partial x_i} \right) \frac{\partial \bar{v}_i}{\partial x_j} + g_i \beta \frac{\nu_t}{\sigma_\theta} \frac{\partial \bar{\theta}}{\partial x_i} \\ &- \varepsilon + \frac{\partial}{\partial x_j} \left[\left(\nu + \frac{\nu_t}{\sigma_k} \right) \frac{\partial k}{\partial x_j} \right] \\ \frac{\partial \varepsilon}{\partial t} + \bar{v}_j \frac{\partial \varepsilon}{\partial x_j} &= \frac{\varepsilon}{k} c_{\varepsilon 1} \nu_t \left(\frac{\partial \bar{v}_i}{\partial x_j} + \frac{\partial \bar{v}_j}{\partial x_i} \right) \frac{\partial \bar{v}_i}{\partial x_j} \\ &+ c_{\varepsilon 1} g_i \frac{\varepsilon}{k} \frac{\nu_t}{\sigma_\theta} \frac{\partial \bar{\theta}}{\partial x_i} - c_{\varepsilon 2} \frac{\varepsilon^2}{k} + \frac{\partial}{\partial x_j} \left[\left(\nu + \frac{\nu_t}{\sigma_\varepsilon} \right) \frac{\partial \varepsilon}{\partial x_j} \right] \end{aligned}$$